arXiv:1202.2267v1 [math.NT] 10 Feb 2012

ON THE SOLUTIONS OF THE EQUATION $(4^n)^x + p^y = z^2$

BILGE PEKER AND SELIN (INAG) CENBERCI

Abstract. In this paper, we gave solutions of the Diophantine equations $16^x + p^y = z^2$, $64^x + p^y = z^2$ where p is an odd prime, $n \in \mathbb{Z}^+$and x, y, z are non-negative integers. Finally we gave a generalization of the Diophantine equation $(4^n)^x + p^y = z^2$.

1. Introduction

There are lots of studies about the Diophantine equation of type $a^x + b^y = c^z$. In 1999, Z. Cao [3] proved that this equation has at most one solution with $z > 1$. In 2005, D. Acu [2] showed that the Diophantine equation $2^x + 5^y = z^2$ has exactly two solutions in non-negative integers, i.e. $(x, y, z) \in \{(3, 0, 3), (2, 1, 3)\}$. J. Sandor [8] studied on the Diophantine equation $4^x + 18^y = 22^z$. In 2011, A. Suvarnamani [11] considered the Diophantine equation $2^x + p^y = z^2$ where p is a prime and x, y, z are non-negative integers. A. Suvarnamani, A. Singta and S. Chotchaisthit [12] found solutions of the Diophantine equations $4^x + 7^y = z^2$ and $4^x + 11^y = z^2$. In 1657, Frenicle de Bessy [7] solved a problem possed by Fermat: if p is an odd prime and $n \geq 2$ is integer, then the equation $x^2 - 1 = p^n$ has no integer solution. S. Chotchaisthit [4] showed all non-negative integer solutions of $4^x + p^y = z^2$ where p is a prime number. B. Peker, S. I. Cenberci [6] studied on the solutions of the Diophantine eqations of some types $(2^n)^x + p^y = z^2$ where p is an odd prime and x, y, z are non-negative integers.

In this study, we gave solutions of the Diophantine equations $16^x + p^y = z^2$, $64^x + p^y = z^2$ where p is an odd prime and x, y, z are non-negative integers. Then we gave a generalization of the Diophantine equation $(4^n)^x + p^y = z^2$.

2. Preliminaries

In this study, we use Catalan's Conjecture [5] and a theorem. Now we give them.

Conjecture 1. *(Catalan) The only solution in integers $a > 1$, $b > 1$, $x > 1$ and $y > 1$ of the equation $a^x - b^y = 1$ is $a = y = 3$ and $b = x = 2$.*

Theorem 1. [7] *If p is an odd prime and $n \geq 2$ is integer, then the equation $x^2 - 1 = p^n$ has no integer solution.*

Now we give our theorems.

Theorem 2. *When p is an odd prime and k, x, y, z are non-negative integers, solutions of the Diophantine equation*

$$16^x + p^y = z^2 \qquad (1)$$

are given in the following two cases:

Date: January 21, 2012.
2000 *Mathematics Subject Classification.* Primary 11D61.
Key words and phrases. Exponential Diophantine Equation.

Case 1. For $p = 3$, $(x, y, z) = (1, 2, 5)$,
Case 2. For $p = 1 + 2^{2k+1}$, $(x, y, z) = \left(k, 1, 2^{2k} + 1\right)$.

Proof. If we consider $y > 0$, then we get

$$z^2 - 4^{2x} = p^y$$

i.e.

$$(z - 4^x)(z + 4^x) = p^y$$

where $z - 4^x = p^v$ and $z + 4^x = p^{y-v}$, $y > 2v$ and v is a non-negative integer. Then we get $p^{y-v} - p^v = 2^{2x+1}$ or $p^v(p^{y-2v} - 1) = 2^{2x+1}$.

If $v = 0$, we obtain $p^y - 1 = 2^{2x+1}$ or $p^y - 2^{2x+1} = 1$. From the *Catalan's Conjecture*, it is obvious that $p = 3$, $y = 2$ and $2x + 1 = 3$, $x = 1$. Hence a solution of the equation (1) is $(x, y, z, p) = (1, 2, 5, 3)$.

For $y = 1$, we obtain $p - 2^{2x+1} = 1$, i.e. $p = 1 + 2^{2x+1}$. Hence solutions of the equation (1) are $(x, y, z) = (x, 1, 2^{2x} + 1)$.

For $y = 0$, we get $z^2 - 4^{2x} = 1$, i.e. $z^2 - 2^{4x} = 1$ which is no solution when $z = 0$ or $z = 1$. From the *Catalan's Conjecture*, $z = 3$ and $4x = 3$ which is impossible.

If we consider $x = 0$, then we have

$$p^y + 1 = z^2$$

i.e.

$$z^2 - p^y = 1.$$

From the *Catalan's Conjecture*, it is obvious that $p = 2$, $y = 3$ and $z = 3$. Hence a solution of the equation (1) is $(x, y, z, p) = (0, 3, 3, 2)$. But this is not a solution, since we assume that p is an odd prime.

Consequently, it is easy to see that $(x, y, z, p) = (1, 2, 5, 3)$ or $(x, y, z, p) = (k, 1, 2^{2k} + 1, 1 + 2^{2k+1})$ where k is a non-negative integer are solutions of the equation $16^x + p^y = z^2$. □

Theorem 3. *When p is an odd prime and k, x, y, z are non-negative integers, solutions of the Diophantine equation*

$$64^x + p^y = z^2 \qquad (2)$$

is given $(x, y, z) = \left(k, 1, 2^{3k} + 1\right)$ *for* $p = 1 + 2^{3k+1}$.

Proof. By using the proof of the *Theorem 2*, solutions of the equation (2) are obvious. □

Now, in the following theorem we will give a generalization of the above Diophantine equations.

Theorem 4. *When p is an odd prime, $n \in \mathbb{Z}^+$ and k, x, y, z are non-negative integers, solutions of the Diophantine equation*

$$(4^n)^x + p^y = z^2 \qquad (3)$$

are given in the following two cases:
Case 1. For $p = 3$, $(x, y, z) = \{(2, 2, 5), (1, 2, 5)\}$,
Case 2. For $p = 1 + 2^{nk+1}$, $(x, y, z) = \left(k, 1, 2^{nk} + 1\right)$.

Proof. If we consider $y > 0$, then we get

$$z^2 - 4^{nx} = p^y$$

i.e.

$$(z - 2^{nx})(z + 2^{nx}) = p^y$$

where $z - 2^{nx} = p^v$ and $z + 2^{nx} = p^{y-v}$, $y > 2v$ and v is a non-negative integer. Then we get $p^{y-v} - p^v = 2^{nx+1}$ or $p^v(p^{y-2v} - 1) = 2^{nx+1}$.

If $v = 0$, we obtain $p^y - 1 = 2^{nx+1}$ or $p^y - 2^{nx+1} = 1$. From the *Catalan's Conjecture*, it is obvious that $p = 3,\ y = 2$ and $nx + 1 = 3,\ nx = 2$. This gives us two cases.

First case: If $n = 1$, then $x = 2$. So we obtain the Diophantine equation $4^x + p^y = z^2$. With this condition, the solution of this equation was given by Chotchaisthit [4] in the form $(x, y, z, p) = (2, 2, 5, 3)$.

Second case: If $n = 2$, then $x = 1$. Therefore we obtain the Diophantine equation $16^x + p^y = z^2$. With this condition, the solution of this equation is given in the *Theorem* 2 in the form $(x, y, z, p) = (1, 2, 5, 3)$.

If we consider $y = 1$, we obtain $p - 2^{nx+1} = 1$, i.e. $p = 1 + 2^{nx+1}$. For $n = 1$, solutions of the equation $4^x + p^y = z^2$ is of the form $(x, y, z, p) = \left(r, 1, 2^r + 1, 2^{r+1} + 1\right)$ where r is a non-negative integer in [4]. For $n = 2$, solutions of the equation (1) is of the form $(x, y, z, p) = \left(k, 1, 2^{2k} + 1, 2^{2k+1} + 1\right)$ where k is a non-negative integer. For $n = 3$, solutions of the equation (2) is given by $(x, y, z, p) = \left(k, 1, 2^{3k} + 1, 2^{3k+1} + 1\right)$ where k is a non-negative integer. If n is any positive integer, solutions of the equation (3) is given of the form $(x, y, z, p) = (k, 1, 2^{nk} + 1, 2^{nk+1} + 1)$ where k is a non-negative integer.

For $y = 0$, we get $z^2 - 4^{nx} = 1$, i.e. $z^2 - 2^{2nx} = 1$ which is no solution when $z = 0$ or $z = 1$. For the other cases if we consider the *Catalan's Conjecture*, we don't find a solution.

If we consider $x = 0$, we have

$$p^y + 1 = z^2$$

i.e.

$$z^2 - p^y = 1.$$

From the *Catalan's Conjecture*, it is obvious that $p = 2,\ y = 3$ and $z = 3$. Hence a solution of the equation (3) is $(x, y, z, p) = (0, 3, 3, 2)$. Since p is not an odd prime, this is not a solution becouse p is an even prime.

Therefore, it is easy to see that $(x, y, z, p) = \{(1, 2, 5, 3), (2, 2, 5, 3)\}$ or $(x, y, z, p) = (k, 1, 2^{nk}+1, 1+2^{nk+1})$ where k is a non-negative integer are solutions of $(4^n)^x + p^y = z^2$. □

References

[1] D. Acu, On the Diophantine equations of type $a^x + b^y = c^z$, General Mathematics Vol.13, No.1, 67-72, 2005.

[2] D. Acu, On a Diophantine equation $2^x + 5^y = z^2$, General Mathematics Vol.15, No.4, 145-148, 2007.

[3] Z. Cao, A note on the Diophantine equation $a^x + b^y = c^z$, Acta Arithmetica XCI, No.1, 85-89, 1999.

[4] S. Chotchaisthit, On the Diophantine equation $4^x + p^y = z^2$ where p is a prime number, American Jr. of Mathematics and Sciences Vol.1, No.1, January 2012.

[5] P. Mihailescu, Primary cyclotomic units and a proof of Catalan's Conjecture, J. Reine Angew Math. 572, 167-195, 2004.
[6] B. Peker, S. I. Cenberci, On the Diophantine equations of $(2^n)^x + p^y = z^2$ type, American. Journal of Mathematics and Sciences Vol.1, No.1, January 2012.
[7] P. Ribenboim, Catalan's Conjecture, Academic Press. Inc. Boston, MA, 1994.
[8] J. Sandor, On a Diophantine equation $4^x + 18^y = 22^z$, Geometric theorems, Diophantine equations and Arithmetic functions, American Research Press Rehobot 4, 91-92, 2002.
[9] W. Sierpinski, Elementary Theory of Numbers, Warszawa, 1964.
[10] J. H. Silverman, A Friendly Introduction to Number Theory, 2nd ed. Prentice- Hall, Inc. New Jersey, 2001.
[11] A. Suvarnamani, Solutions of theDiophantine equation $2^x + p^y = z^2$, Int. J. of Mathematical Sciences and Applications Vol.1, No.3, September 2011.
[12] A. Suvarnamani, A. Singta and S. Chotchaisthit, On two Diophantine equations $4^x + 7^y = z^2$ and $4^x + 11^y = z^2$, Science and Technology RMUTT Journal Vol.1, No.1, 2011.

Elementary Mathematics Education Programme, Ahmet Kelesoglu Education Faculty, Konya University, KONYA, TURKEY.
E-mail address: bilge.peker@yahoo.com

Secondary Mathematics Education Programme, Ahmet Kelesoglu Education Faculty, Konya University,KONYA, TURKEY.
E-mail address: inag_s@hotmail.com